# Genetic analysis of individual breeds of poultry used in breeding programs in Slovakia











Cyril Hrnčár

Slovak University of Agriculture in Nitra

Faculty of Agrobiology and Food Resources



- the current economic situation causes an interest in specialized poultry lines in advanced countries in the world, focusing on egg-laying or meat-based performance,
- due to the very intense breeding, many genes are lost and they may be missing in future,
- therefore, the reasons for keeping poultry genetic resources are the same as for all species of farm animals.



- at present, with the growing demand for poultry products from extensive systems, it is important to raise native poultry breeds suitable for free-range and organic farming because of their good adaptation to the local conditions,
- native poultry breeds play an important social role and have a positive impact on maintaining rural society and traditional form of agriculture as well as gratify local traditions,
- they represent a gene source for future breeding strategies and research,
- an alternative native breeds in poultry production system have an important advantage of biodiversity and sustainable agricultural production.



# **AUTHOCHTONOUS POULTRY BREEDS**

chicken breeds (Oravka, Bantam Oravka),

# goose breeds (Slovak Goose, Suchovska Goose).



- dual-purpose poultry breed with good egg production, growth ability and adaptability on alternative outdoor,
- the goal was to develop a breed suitable for harsh climatic conditions of northern Slovakia which can be kept in free range,
- hard rectangular frame, comb is rosette, legs are medium length without feather with yellow color with red tinted strip (lampas), feather is rich with solid structure.



# **PERFORMANCE OF ORAVKA**

- body weight: cock 2.8-3.3 kg, hen 2-2.-2.7 kg,
  - ♣ laying: 180-200 eggs,
- 🎍 🖶 egg weight: 58 g,
  - brownish eggshell.





#### YELLOW-BROWNISH ORAVKA

- the oldest and most common color variety of Oravka,
- created by combinatorial crossing of regional breeds with breeds of Rhode Island Red, Wyandotte White and New Hampshire,
- recognised in year 1990.








# **WHITE ORAVKA**

- created by crossing of Yellow-brownish Oravka with Plymouth Rock White,
  - recognised in year 2008.







#### **BLACK ORAVKA**

- created by crossing of White Oravka with breeds of Dresdner Black, Plymouth Rock Black and Barnevelder Black,
- recognised in year 2017.







#### **CRELE ORAVKA**

- a combination of rusty partridge colorations with sparrowhawk pattern,
- created by crossing of Yellow-brawnish Oravka, White Oravka and Crele Bielefelder,







#### **BARRED ORAVKA**

- created by crossing of White Oravka with Barred American Rock,







#### **CURRENT STATUS OF ORAVKA**

- Slovak University of Agriculture in Nitra:
  - ex situ conservation flock of Yellow-brownish Oravka,
  - monitoring of population size of Oravka in Slovakia,
  - performance (ejaculate quality, hatching, growth ability, egg production, egg quality, carcass parameters, meat quality),
  - creation of the system and method of authentication for reliably distinguish between the Oravka and other chicken breeds by analyse chicken feathers (use 7 microsatellite markers recommended by FAO) - DAPC analysis.



#### **CURRENT STATUS OF ORAVKA**

- National Agriculture and Food Centre –Research Institute for Animal Production (NAFC–RIAP) Nitra:
  - ex situ conservation flock of Yellow brownish Oravka,
  - performance (hatching, growth ability, egg production, egg quality)
  - isolation, culture and cryopreservation of blastodermal and primordial germ cells in Oravka chickens,
  - the quality of blastodermal and primordial germ cells, before and after cryopreservation is evaluated by fluorescent microscopy, flow cytometry and transmission electron microscopy,
  - o cryopreserve roosters semen of Oravka breed and assess their quality by methods mentioned above, semen samples of individual roosters are stored in the gene bank at NAFC-RIAP Nitra.



#### **CURRENT STATUS OF ORAVKA**

- Slovak Association of Breeders:
  - creation of new color variants of Oravka in reproductive flocks with individual laying and pedigree hatching,
  - o exterior judging of Oravka at poultry exhibitions and recognizing of reproductive flocks,
  - o monitoring of laying (collective, individual) of Oravka hens in reproductive flocks.



#### **BANTAM ORAVKA**

- bantam form of large breed,
- ♣ Bantam Oravka should follow the large fowl standard in all respects, however, the feathering color is more intense compared with large Oravka,
- dual-purpose poultry breed with good egg production, growth ability and adaptability on alternative outdoor.





#### PERFORMANCE OF BANTAM ORAVKA

- ♣ body weight: cock 1.0-1.3 kg, hen 0.9-1.1 kg,
- 4 laying: 140 eggs,
- egg weight: 40 g,
- brownish eggshell.





# YELLOW-BROWNISH BANTAM ORAVKA

- breed was developed in the 1980s as the result of crossbreeding of large Oravka with Bantam Sumavian and Bantam Dresdner,
- recognised in year 1990.







# WHITE BANTAM ORAVKA

- created by crossing of Yellow-brownish Bantam Oravka, White Bantam Dresdner and White Bantam Plymouth Rock,
- recognised in year 2017.







# **CRELE BANTAM ORAVKA**

- a combination of rusty partridge colorations with sparrowhawk pattern,
- created by crossing of Yellow-brownish Bantam Oravka and Crele Bantam Bielefelder,
- recognised in year 2017.







# **CURRENT STATUS OF BANTAM ORAVKA**

- Slovak University of Agriculture in Nitra:
  - performance (egg production, egg quality).



#### **CURRENT STATUS OF BANTAM ORAVKA**

#### Slovak Association of Breeders:

- creation of new color variants of Bantam Oravka in reproductive flocks with individual laying and pedigree hatching,
- exterior judging of Bantam Oravka at poultry exhibitions and recognizing of reproductive flocks,
- o monitoring of laying (collective, individual) of Bantam Oravka in reproductive flocks.



- established in 1940s on the basis of regional breeds from South-Eastern part of Slovakia (from Nitra and Levice areas) by crossing with goose breed from Germany and Hungary,
- the aim of breeding was to create a medium weight triple purpose (meat, liver, feather) goose suitable for corn areas, a strong,
- resistant goose with a good pasturing ability and with preserved clucking instinct,
- first standard of the Slovak goose was published by Malík (1966),







#### **SLOVAK GOOSE**

- middle-heavy breed of goose noble appearance and firm constitution. It uses very good pasture and has preserved quiltiness,
- feather is pure white, made of medium quality feathers, the volume fraction of the punch exceeds the volume of the cover feather,
- hardy, good for grazing. It has a preserved quandary instinct that is itself hatch and inferred by the young,
- good disposition for cooking and creating delicious livers.







# PERFORMANCE OF SLOVAK GOOSE

- y → body weight: gender 6.8-8.0 kg, goose 5.7-7.0 kg,
  - laying: 12 and more eggs,
  - egg weight: 140 g,
  - white eggshell.







#### **SUCHOVSKA GOOSE**

- established at the end of the 1980s in the village Suchá nad Parnou from the population of local yellow coloured domestic goose by cross-breeding with French (Toulouse, Landes) and German (Pomeranian, Steinbacher) breeds of geese,
- medium-heavy triple purpose geese, preservation of clucking instinct,
- created to breed goose of bigger body frame, firm constitution and of compact and solid body,
- recognized in year 1995.







#### **SUCHOVSKA GOOSE**

- bigger and hard body frame, te body is long, compact, muscled, firm, almost horizontally held, body width is equal to its height,
- saddling and pasture type,
- 4 feather is smoother, with more dust, color is yellow-wild,
- female is similarly colored as a male with a larger amount of gray admixture, overall appears darker.







#### PERFORMANCE OF SUCHOVSKA GOOSE

- - ♣ laying: 12-16 eggs,
  - egg weight: 140 g,
  - white eggshell.







#### **CURRENT STATUS OF SLOVAK AND SUCHOVSKA GOOSE**

- Slovak University of Agriculture in Nitra:
  - monitoring of population size of Slovak and Suchovska goose in Slovakia,
  - performance (hatching, growth ability, carcass parameters, meat quality).
  - histological and histochemical evalution of geese muscle tissue,
  - o characterization of the genetic diversity and structure of both native Slovak goose breeds based on six microsatellite loci analysis.



# **CURRENT STATUS OF SLOVAK AND SUCHOVSKA GOOSE**

- National Agriculture and Food Centre –Research Institute for Animal Production (NAFC–RIAP) Nitra:
  - cryopreserve gander semen of Slovak and Suchovska Goose breeds,
  - o semen samples of individual ganders are stored in the gene bank at NPPC-RIAP Nitra.



# **CURRENT STATUS OF SLOVAK AND SUCHOVSKA GOOSE**

- Slovak Association of Breeders:
  - exterior judging of Slovak and Suchovska Goose at poultry exhibitions and recognizing of reproductive flocks.



#### **FUTURE BREEDING GOALS**

- focusing on the improvement of exterior characteristics of native chickens a goose breeds,
- the small number of recognized flocks puts increased pressure on poultryjudges when judging poultry at shows and recognizing of reproductive flocks,
- cooperetion the Slovak Association of Breeders with SUA in Nitra and NAFC-RIAP Nitra for increasing performance of Oravka (ejaculate quality, egg production, egg quality, growt ability, meat quality),
- a high risk for the reproduction of Slovak Goose is a tendency of gander, male goose, to be monogamic and the mutual refusal, which exist in both sexes.



# **Partners:**





# Thank you for your attention!

This presentation has been supported by the Erasmus+ KA2 Cooperation Partnerships grant no. 2021-1-SK01-KA220-HED-000032068 "Innovation of the structure and content of study programs in the field of animal genetic and food resources management with the use of digitalisation - Inovácia obsahu a štruktúry študijných programov v oblasti manažmentu živočíšnych genetických a potravinových zdrojov s využitím digitalizácie". The European Commission support for the production of this presentation does not constitute an endorsement of the contents which reflects the views only of the authors, and the Commission cannot be held responsible for any use which may be made of the information contained therein.



Cyril Hrnčár



cyril.hrncar@uniag.sk



